

Dear Author

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For **fax** submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style.
- Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL:

<http://dx.doi.org/10.1007/s11302-013-9403-2>

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information, go to:

<http://www.springerlink.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us, if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

| | | | |
|----|--------------------------|--|-------------------------------|
| 1 | Article Title | Investigation of the functional expression of purine and pyrimidine receptors in porcine-isolated pancreatic arteries | |
| 2 | Article Sub- Title | | |
| 3 | Article Copyright - Year | Springer Science+Business Media Dordrecht 2013 (This will be the copyright line in the final PDF) | |
| 4 | Journal Name | Purinergic Signalling | |
| 5 | Corresponding Author | Family Name | Ralevic |
| 6 | | Particle | |
| 7 | | Given Name | V. |
| 8 | | Suffix | |
| 9 | Author | Organization | University of Nottingham |
| 10 | | Division | School of Biomedical Sciences |
| 11 | | Address | Nottingham NG7 2UH, UK |
| 12 | | e-mail | vera.ralevic@nottingham.ac.uk |
| 13 | Author | Family Name | Alsaqati |
| 14 | | Particle | |
| 15 | | Given Name | M. |
| 16 | | Suffix | |
| 17 | Author | Organization | University of Nottingham |
| 18 | | Division | School of Biomedical Sciences |
| 19 | | Address | Nottingham NG7 2UH, UK |
| 20 | | e-mail | |
| 21 | Author | Family Name | Chan |
| 22 | | Particle | |
| 23 | | Given Name | S. L. F. |
| 24 | | Suffix | |
| 25 | Author | Organization | University of Nottingham |
| 26 | | Division | School of Biomedical Sciences |
| 27 | | Address | Nottingham NG7 2UH, UK |
| 28 | | e-mail | |
| 29 | Schedule | Received | 2 July 2013 |
| 30 | | Revised | |

| | | | |
|----|-----------------------------|--|-----------------|
| 31 | | Accepted | 5 November 2013 |
| 32 | Abstract | <p>Receptors for purines and pyrimidines are expressed throughout the cardiovascular system. This study investigated their functional expression in porcine-isolated pancreatic arteries. Pancreatic arteries (endothelium intact or denuded) were prepared for isometric tension recording and preconstricted with U46619, a thromboxane A₂ mimetic; adenosine-5'-diphosphate (ADP), uridine-5'-triphosphate (UTP) and MRS2768, a selective P₂Y₂ agonist, were applied cumulatively, while adenosine-5'-triphosphate (ATP) and $\alpha\beta$-methylene-ATP ($\alpha\beta$-meATP) response curves were generated from single concentrations per tissue segment. Antagonists/enzyme inhibitors were applied prior to U46619 addition. ATP, $\alpha\beta$-meATP, UTP and MRS2768 induced vasoconstriction, with a potency order of $\alpha\beta$-meATP > MRS2768 > ATP \geq UTP. Contractions to ATP and $\alpha\beta$-meATP were blocked by NF449, a selective P₂X₁ receptor antagonist. The contraction induced by ATP, but not UTP, was followed by vasorelaxation. Endothelium removal and DUP 697, a cyclooxygenase-2 inhibitor, had no significant effect on contraction to ATP but attenuated that to UTP, indicating actions at distinct receptors. MRS2578, a selective P₂Y₆ receptor antagonist, had no effect on contractions to UTP. ADP induced endothelium-dependent vasorelaxation which was inhibited by MRS2179, a selective P₂Y₁ receptor antagonist, or SCH58261, a selective adenosine A_{2A} receptor antagonist. The contractions to ATP and $\alpha\beta$-meATP were attributed to actions at P₂X₁ receptors on the vascular smooth muscle, whereas it was shown for the first time that UTP induced an endothelium-dependent vasoconstriction which may involve P₂Y₂ and/or P₂Y₄ receptors. The relaxation induced by ADP is mediated by P₂Y₁ and A_{2A} adenosine receptors. Porcine pancreatic arteries appear to lack vasorelaxant P₂Y₂ and P₂Y₄ receptors.</p> | |
| 33 | Keywords separated by ' - ' | <p>$\alpha\beta$-meATP - ATP - UTP - ADP - MRS2578 - P₂Y₁ - P₂Y₂ - P₂X₁ - A_{2A} adenosine receptors - Vasoconstriction - Relaxation - Endothelium</p> | |
| 34 | Foot note information | | |

Investigation of the functional expression of purine and pyrimidine receptors in porcine-isolated pancreatic arteries

M. Alsaqati · S. L. F. Chan · V. Ralevic

Received: 2 July 2013 / Accepted: 5 November 2013
© Springer Science+Business Media Dordrecht 2013

Abstract Receptors for purines and pyrimidines are expressed throughout the cardiovascular system. This study investigated their functional expression in porcine-isolated pancreatic arteries. Pancreatic arteries (endothelium intact or denuded) were prepared for isometric tension recording and preconstricted with U46619, a thromboxane A₂ mimetic; adenosine-5'-diphosphate (ADP), uridine-5'-triphosphate (UTP) and MRS2768, a selective P2Y₂ agonist, were applied cumulatively, while adenosine-5'-triphosphate (ATP) and $\alpha\beta$ -methylene-ATP ($\alpha\beta$ -meATP) response curves were generated from single concentrations per tissue segment. Antagonists/enzyme inhibitors were applied prior to U46619 addition. ATP, $\alpha\beta$ -meATP, UTP and MRS2768 induced vasoconstriction, with a potency order of $\alpha\beta$ -meATP > MRS2768 > ATP \geq UTP. Contractions to ATP and $\alpha\beta$ -meATP were blocked by NF449, a selective P2X₁ receptor antagonist. The contraction induced by ATP, but not UTP, was followed by vasorelaxation. Endothelium removal and DUP 697, a cyclooxygenase-2 inhibitor, had no significant effect on contraction to ATP but attenuated that to UTP, indicating actions at distinct receptors. MRS2578, a selective P2Y₆ receptor antagonist, had no effect on contractions to UTP. ADP induced endothelium-dependent vasorelaxation which was inhibited by MRS2179, a selective P2Y₁ receptor antagonist, or SCH58261, a selective adenosine A_{2A} receptor antagonist. The contractions to ATP and $\alpha\beta$ -meATP were attributed to actions at P2X₁ receptors on the vascular smooth muscle, whereas it was shown for the first time that UTP induced an endothelium-dependent vasoconstriction which may involve P2Y₂ and/or P2Y₄ receptors. The relaxation induced by ADP is mediated by P2Y₁ and A_{2A}

adenosine receptors. Porcine pancreatic arteries appear to lack vasorelaxant P2Y₂ and P2Y₄ receptors.

Keywords $\alpha\beta$ -meATP · ATP · UTP · ADP · MRS2578 · P2Y₁ · P2Y₂ · P2X₁ · A_{2A} adenosine receptors · Vasoconstriction · Relaxation · Endothelium

Abbreviations

| | |
|----------------------|--|
| $\alpha\beta$ -meATP | $\alpha\beta$ -Methylene-adenosine-5'-triphosphate |
| ADP | Adenosine-5'-diphosphate |
| ATP | Adenosine-5'-triphosphate |
| EDCFs | Endothelium-derived contractile factors |
| ENTPDase | Ecto-nucleotidase 5'-triphosphate diphosphohydrolase |
| PPADS | Pyridoxal phosphate-6-azo(benzene-2,4-disulfonic acid) |
| UTP | Uridine-5'-triphosphate |
| VSMCs | Vascular smooth muscle cells |
| XAC | Xanthine amine congener |

Introduction

The activities of both exocrine and endocrine cells of the pancreas are regulated by autonomic nerves (parasympathetic and sympathetic) as well as by hormones and autocrine and paracrine mediators. Although the exact mechanisms remain to be established, it is generally agreed that an increase in endocrine cell activity during hormone secretion corresponds with an increase in blood flow, to meet the metabolic demand. The role of exogenous purine and pyrimidine nucleotides in controlling the functions of endocrine and exocrine components of the pancreas are well described [1, 2], but little is known about their effects on pancreatic arterial vasocontractility.

M. Alsaqati · S. L. F. Chan · V. Ralevic (✉)
School of Biomedical Sciences, University of Nottingham,
Nottingham NG7 2UH, UK
e-mail: vera.ralevic@nottingham.ac.uk

84 There are two main families of P2 purine and pyrimidine
 85 receptors, ionotropic P2X and G protein-coupled P2Y recep-
 86 tors. Molecular cloning has identified seven mammalian P2X
 87 receptor subunits: P2X1, P2X2, P2X3, P2X4, P2X5, P2X6
 88 and P2X7 [3], while eight mammalian P2Y receptors have
 89 been identified: P2Y₁, P2Y₂, P2Y₄, P2Y₆, P2Y₁₁, P2Y₁₂,
 90 P2Y₁₃ and P2Y₁₄ [4]. P2X receptors are activated by
 91 adenosine-5'-triphosphate (ATP) and its stable, and conse-
 92 quently more potent, analogue $\alpha\beta$ -methylene-ATP ($\alpha\beta$ -
 93 meATP) [5, 6]. P2Y receptors can be divided on the basis of
 94 their endogenous agonists into adenine nucleotide-preferring
 95 (P2Y₁, P2Y₁₁, P2Y₁₂ and P2Y₁₃) receptors and uracil nucle-
 96 otide or UDP-sugar-preferring (P2Y₂, P2Y₄, P2Y₆ and
 97 P2Y₁₄) receptors [7]. Among the adenine nucleotide group,
 98 the human P2Y₁₁ receptor is selectively activated by ATP and
 99 fails to respond to adenosine-5'-diphosphate (ADP) [8], al-
 100 though the dog orthologue responds to both ADP and ATP
 101 [9]. P2Y₁, P2Y₁₂ and P2Y₁₃ receptors are activated by ADP,
 102 with lower potency by ATP [10–13]. Among the uracil nucle-
 103 otide or UDP-sugar receptors, P2Y₂ is equally activated by
 104 ATP and uridine-5'-triphosphate (UTP), while P2Y₄ receptor
 105 is highly selective for UTP over ATP [14]. The P2Y₆ receptor
 106 is activated by UDP and UTP, while the P2Y₁₄ receptor is
 107 activated by UDP and UDP-sugars [6, 15].
 108 Within the pancreatic vasculature, P2X1, P2X2, P2Y₁ and
 109 P2Y₂ receptors were detected by immunohistochemistry [16].
 110 More than two decades ago, it was shown that P2X receptors
 111 mediate pancreatic vasoconstriction, and P2Y receptors me-
 112 diate vasodilatation in response to ATP [17], and subsequent
 113 studies showed an additional involvement of contractile recep-
 114 tors sensitive to UTP (named P2U receptors) [18]. Purine
 115 receptor subclassification has advanced significantly since
 116 that time. A re-evaluation of purine receptors in the pancreatic
 117 vasculature is clearly warranted. In the current study, we
 118 describe the pharmacological characterisation of P2Y₁ and
 119 A_{2A} receptor-mediated relaxatory responses, in addition to
 120 P2X1, P2Y₂ and P2Y₄ receptor-mediated contractile re-
 121 sponses of porcine-isolated pancreatic artery preparations.
 122 P2Y₂ and/or P2Y₄ receptors appear to be expressed mainly
 123 in endothelial cells, while P2X1 and A_{2A} receptors appear to
 124 be expressed in smooth muscle cells of the pancreatic arteries.
 125 A preliminary account of some of these data has previously
 126 been presented to the British Pharmacological Society [19].

127 Materials and methods

128 Tissue preparation

129 Pancreases from pigs (either sex, age less than 6 months, wt
 130 ~50 kg) were obtained on ice from a local abattoir (G Wood &
 131 Sons Ltd., Mansfield). A crude dissection was conducted to
 132 isolate the porcine pancreatic arteries (greater pancreatic

133 artery) which were located in the body of the pancreas. The
 134 vessels were dissected out and placed in Krebs–Henseleit
 135 buffer containing 2 % (*w/v*) Ficoll (hydrophilic polysaccha-
 136 ride, type 70) and were refrigerated overnight at 4 °C. The
 137 next day, a fine dissection was performed on arteries, and the
 138 artery segments were cut into rings of about 0.5 cm in length
 139 and suspended in Krebs–Henseleit buffer (gassed, 95 % O₂,
 140 5 % CO₂).

141 The endothelium of some arteries was removed by gently
 142 rubbing the innermost surface of the artery with forceps on a
 143 paper tissue before attaching it to the set-up [20]. Successful
 144 removal of the endothelium was tested using substance P
 145 (10 nM). Endothelium-denuded arteries relaxed in response
 146 to substance P to less than 10 % of the U46619-induced
 147 contraction, while in endothelium-intact arteries, the relaxa-
 148 tion to substance P was 36 %±8 (*n*=7, data not shown).

Responses in the porcine-isolated pancreatic artery 149

150 Arterial rings were mounted onto wires in tissue baths con-
 151 taining warm (37 °C), oxygenated Krebs–Henseleit solution
 152 and were connected via isometric force transducers
 153 (mechanotransducer MLT 050/D; ADInstruments, Sydney,
 154 Australia) to a PC running the computer programme,
 155 LabChart (ADInstruments, Sydney, Australia). Rings were
 156 put under tension (15 g) and allowed to equilibrate for
 157 60 min before assessing the viability with two challenges of
 158 75 mM potassium chloride (KCl). The tissues were then
 159 allowed to equilibrate for 60 min, after which U46619 (10–
 160 100 nM), a thromboxane A₂ mimetic, was used to contract the
 161 tissues to between 40 and 80 % of the second KCl response.
 162 This ensured that if there was a vasodilator component to the
 163 response, this could be detected. Once an appropriate level of
 164 U46619 response had been achieved, ATP, $\alpha\beta$ -meATP, UTP,
 165 ADP or MRS2768 were added. Antagonists or enzyme inhib-
 166 itors were applied 10 min prior to the addition of U46619,
 167 allowing them to be incubated with the tissues for a minimal
 168 contact time of 30 min prior to the application of agonists.
 169 Some arteries were incubated with 0.1 % (*v/v*) DMSO (vehi-
 170 cle control).

Reagents and drugs 171

172 Krebs–Henseleit buffer was composed of the following (mM):
 173 NaCl, 118; KCl, 4.8; CaCl₂·H₂O, 1.3; NaHCO₃, 25.0;
 174 KH₂PO₄, 1.2; MgSO₄·7H₂O, 1.2; and glucose, 11.1. Suramin,
 175 UTP, ATP, $\alpha\beta$ -meATP, ADP, U46619, xanthine amine con-
 176 gener (XAC) and SCH58261 (7-(2-phenylethyl)-5-amino-
 177 2-(2-furyl)-pyrazolo-[4,3-*e*]-1,2,4-triazolo[1,5-*c*]pyrimidine)
 178 were purchased from Sigma (Poole, Dorset, UK), while DUP
 179 697 (5-bromo-2-(4-fluorophenyl)-
 180 3-[4-(methanesulfonyl)phenyl]-thiophene), pyridoxal
 181 phosphate-6-azo(benzene-2,4-disulfonic acid) (PPADS),

MRS2578 (*N,N'*-1,4-butanediyl bis(*N'*-[3-isothiocyanatophenyl] thiourea)), MRS2179 (2'-deoxy-*N*⁶-methyladenosine 3',5'-bisphosphate tetrasodium salt), MRS2768 (uridine-5'-tetrphosphate δ -phenyl ester tetrasodium salt) and substance P were purchased from Tocris Biosciences Ltd. (Bristol, UK). NF449 (4,4',4'',4'''-[carbonylbis(imino-5,1,3-benzenetriyl-bis(carbonylimino))] tetrakis-1,3-benzenedisulfonic acid) was purchased from Calbiochem-Merck4Biosciences. U46619 was dissolved in ethanol at 10 mM stock concentration. PPADS, suramin, $\alpha\beta$ -meATP, ATP, ADP, UTP, NF449, MRS2179, MRS2768 and substance P were dissolved in distilled water. DUP 697, XAC, MRS2578 and SCH58261 were dissolved in DMSO at 10 mM stock concentration.

Data analysis

The contractions to ATP, $\alpha\beta$ -meATP and UTP were measured from the stabilised U46619-induced response and were expressed in grams, while the relaxations to ATP and ADP were expressed as a percentage of the U46619-induced contraction. Data were expressed as log concentration–response plots. Values for all figures refer to mean \pm standard error of the mean (SEM) with 95 % confidence. Results were compared by one- or two-way ANOVA with Bonferroni's post hoc test or unpaired Student's *t* test (Prism, GraphPad, San Diego, CA, USA). Differences were considered to be significant when the *P* value was <0.05. *N* expresses the number of animals.

Results

Effect of purine and pyrimidine nucleotides on vascular tone in porcine-isolated pancreatic arteries

To investigate the effect of purine and pyrimidine nucleotide agonists on porcine pancreatic arteries, $\alpha\beta$ -meATP (10 nM to 100 μ M), ATP (1 μ M to 10 mM), UTP (10 μ M to 1 mM), ADP (1 μ M to 1 mM) and MRS2768 (100 nM to 30 μ M) were applied after precontraction with U46619. The responses to ATP and $\alpha\beta$ -meATP were found to be desensitised rapidly. Therefore, they were applied at single concentrations (one concentration per tissue segment). The responses to UTP, ADP and MRS2768 did not desensitise rapidly; thus, cumulative concentration–response curves were generated. ATP, $\alpha\beta$ -meATP, UTP and MRS2768 induced concentration-dependent contraction with a potency order of $\alpha\beta$ -meATP > MRS2768 > ATP \geq UTP (*P* < 0.001, two-way ANOVA; Fig. 1a). The response to ATP was biphasic, since its contraction was followed by a relaxation (Fig. 1b) which was equipotent to the concentration-dependent relaxation produced by ADP (Fig. 1a). The efficacies of ATP and $\alpha\beta$ -




Fig. 1 **a** Concentration-dependent contraction of ATP, $\alpha\beta$ -meATP, UTP and MRS2768, a selective P2Y₂ agonist, and relaxation of ADP and ATP in U46619-precontracted porcine pancreatic arteries (*n* = 7–12). **b** Typical trace showing the biphasic response to ATP (contraction followed by relaxation). Data are presented as mean \pm SEM

meATP in inducing contraction were similar and greater than those of UTP or MRS2768. The relaxation to ADP and ATP at the highest concentration of the agonists used (1 mM) was similar at 4.5 ± 0.5 g (*n* = 10) and 5.5 ± 0.2 g (*n* = 7), respectively; there was no significant difference between these responses (Fig. 1a). UTP, MRS2768 and $\alpha\beta$ -meATP did not elicit vasorelaxation.

Characterisation of responses to ATP and $\alpha\beta$ -meATP in U46619-precontracted porcine-isolated pancreatic arteries

- Effect of suramin, PPADS and $\alpha\beta$ -meATP
- Responses to ATP and $\alpha\beta$ -meATP were characterised using the non-selective P2 receptor antagonists, suramin (100 μ M) and PPADS (10 μ M). Both suramin and PPADS significantly attenuated the contractions evoked by ATP (1 mM) and $\alpha\beta$ -meATP (1 μ M) (Fig. 2a, b). These concentrations of ATP and $\alpha\beta$ -meATP were chosen since they produced robust and submaximal responses, and for $\alpha\beta$ -meATP, the concentration was close to the half maximal effective concentration (EC₅₀) value [mean EC₅₀ value was 1.6 μ M (95 % confidence interval (CI), 1.05 to 2.53 μ M; *n* = 8; Fig. 1a)]. The relaxation to ATP was not affected by suramin or PPADS (Fig. 2c). Since $\alpha\beta$ -meATP induces desensitisation of P2X receptors more

readily than ATP because it is broken down more slowly than ATP [5], the responses to ATP and $\alpha\beta$ -meATP were studied in the presence of $\alpha\beta$ -meATP, in which $\alpha\beta$ -meATP (1 μ M) was added 10 min prior the addition of U46619. As seen in Fig. 2a, b, the contractions to ATP and $\alpha\beta$ -meATP were reduced in the presence of the




Fig. 2 Effect of suramin (100 μ M), PPADS (10 μ M) and desensitisation by $\alpha\beta$ -meATP (1 μ M) on contractions to **a** ATP (1 mM) and **b** $\alpha\beta$ -meATP (1 μ M) and **c** on the relaxation to ATP in U46619-precontracted porcine pancreatic arteries. PPADS, suramin and $\alpha\beta$ -meATP reduced the contractions of **a** ATP and **b** $\alpha\beta$ -meATP (** P <0.01; *** P <0.001, one-way ANOVA with Bonferroni's post hoc test, responses of ATP or $\alpha\beta$ -meATP vs their responses in the presence of PPADS, suramin or $\alpha\beta$ -meATP, n =6–9). **c** The relaxation to ATP was not significantly different in the absence or presence of PPADS, suramin or $\alpha\beta$ -meATP (n =7). Data are presented as mean \pm SEM

desensitising agent, while the relaxation to ATP was not affected (Fig. 2c).

- Effect of NF449, a selective P2X1 receptor antagonist
- Contractile responses to $\alpha\beta$ -meATP suggest an expression of P2X1 receptors in porcine pancreatic arteries (Fig. 2b). In turn, the involvement of P2X1 receptors in contraction to ATP seems likely because contraction was significantly blocked by $\alpha\beta$ -meATP (Fig. 2a). The responses to ATP and $\alpha\beta$ -meATP were studied further in the presence of NF449 (10 μ M), a P2X1 receptor-selective antagonist. The contractions to ATP and $\alpha\beta$ -meATP were inhibited in the presence of NF449 (Fig. 3).

- Effect of endothelium removal
- The response to ATP was tested after the endothelium had been removed. The contraction and the relaxation induced by ATP (Fig. 4) were statistically not significantly different in the absence or presence of the endothelium. Similarly, removal of the endothelium had no effects on the contractions to KCl, U46619 or $\alpha\beta$ -meATP; for example, the contraction to 75 mM KCl was 9.5 ± 0.5 g in endothelium-intact arteries, while it was 9 ± 0.5 g in endothelium-denuded arteries (n =7–9). The contraction




Fig. 3 Effect of NF449 (10 μ M), a selective P2X1 receptor antagonist, on contractions to **a** ATP (1 mM) and **b** $\alpha\beta$ -meATP (1 μ M), in U46619-precontracted porcine pancreatic arteries. NF449 reduced the effects of **a** ATP and **b** $\alpha\beta$ -meATP (** P <0.001, unpaired Student's t test, n =10–13). Data are presented as mean \pm SEM




Fig. 4 Effect of removal of the endothelium on **a** contraction and **b** relaxation to ATP (1 mM) in U46619-precontracted porcine pancreatic arteries. The effect of the removal of endothelium on the contraction or relaxation of ATP was not significantly different ($n=9-11$). Data are presented as mean \pm SEM




Fig. 5 Effect of XAC (10 μ M) on **a** contraction and **b** relaxation to ATP (1 mM) in U46619-precontracted porcine pancreatic arteries. **a** XAC had no effect on the contraction to ATP ($n=8-10$), and **b** XAC reduced the relaxation to ATP ($***P<0.001$, unpaired Student's t test, $n=8-10$). Data are presented as mean \pm SEM

to 10–100 nM U46619 was 5.5 ± 0.5 g in endothelium-intact arteries, while it was 5.8 ± 0.6 g in endothelium-denuded arteries ($n=12-14$). The contraction to 1 μ M $\alpha\beta$ -meATP was 3.2 ± 0.6 g in endothelium-intact arteries, while it was 3 ± 0.6 g in endothelium-denuded arteries ($n=6$); there was no significant difference between these responses.

- Effect of XAC, an adenosine receptor antagonist
The relaxation to ATP was investigated in the presence of a non-selective adenosine receptor antagonist; XAC (10 μ M) had no effect on the contraction evoked by ATP (Fig. 5a), while it reduced significantly the relaxation to ATP (Fig. 5b).

Characterisation of response to UTP in U46619-precontracted porcine-isolated pancreatic arteries

- Effect of suramin, PPADS, $\alpha\beta$ -meATP and MRS2578, a selective P2Y₆ receptor antagonist
The contraction to UTP was examined in the presence of suramin (100 μ M), PPADS (10 μ M), $\alpha\beta$ -meATP

(1 μ M) and MRS2578 (10 μ M). Suramin and PPADS significantly reduced the contraction to UTP (Fig. 6), while the UTP responses were not affected after P2X receptor desensitisation in the presence of $\alpha\beta$ -meATP (1 μ M) or in the presence of a selective P2Y₆ receptor antagonist (MRS2578); for example, the contraction to 1 mM UTP was 1.8 ± 0.2 g in the absence of MRS2578 ($n=7$), while it was 2.1 ± 0.2 g in the presence of




Fig. 6 Effect of suramin (100 μ M) and PPADS (10 μ M) on contraction to UTP in U46619-precontracted porcine pancreatic arteries. With suramin and PPADS, effect of UTP concentration ($F=16.77$ and $F=12.38$, respectively, $***P<0.001$), suramin and PPADS reduced the contraction evoked by UTP ($F=14.47$ and $F=12.48$, respectively, $***P<0.001$, two-way ANOVA; $n=9-12$). Data are presented as mean \pm SEM

MRS2578 ($n=6$); there was no significant difference between these responses.

Effect of endothelium removal

The effects of UTP were studied after the endothelium had been removed. The contraction induced by UTP was significantly attenuated in the endothelium-denuded arteries (Fig. 7).

Effect of DUP 697, a cyclooxygenase-2 inhibitor

Because the contraction to UTP was largely endothelium-dependent, the contraction was studied in the presence of DUP 697, a cyclooxygenase-2 (COX-2) inhibitor, since COX-2 facilitates the release of agents which are responsible for endothelium-dependent contraction. DUP 697 (3 μ M) diminished the response to UTP (Fig. 8) to a similar extent as removal of the endothelium (Fig. 7), while DUP 697 did not alter the contraction to U46619 (the precontraction agent) or the contraction to ATP (data not shown).




Fig. 8 Effect of DUP 697 (3 μ M), a cyclooxygenase-2 inhibitor, on contraction to UTP in U46619-precontracted porcine pancreatic arteries. Effect of UTP concentration ($F=8.48$, $***P<0.001$) and DUP 697 reduced the contraction evoked by UTP ($F=50.8$, $***P<0.001$, two-way ANOVA; $n=8-12$). Data are presented as mean \pm SEM

reduced in the presence of this inhibitor which indicates the involvement of adenosine receptors (Fig. 10). To find out about the adenosine subtype involved in the relaxation to ADP, the response to ADP was investigated in the presence of SCH58261, a selective adenosine A_{2A} receptor antagonist. This antagonist significantly inhibited the relaxation to ADP, to a similar extent as seen with XAC

Characterisation of response to ADP in U46619-precontracted porcine-isolated pancreatic arteries

Effect of MRS2179, a $P2Y_1$ receptor selective antagonist, and of endothelium removal

The relaxation to ADP in pancreatic arteries was studied in the presence of MRS2179 (10 μ M) and after the endothelium had been removed. The relaxation to ADP was reduced slightly but significantly in the presence of MRS2179 (Fig. 9a) and in the endothelium-denuded arteries (Fig. 9b), which indicates the involvement of $P2Y_1$ receptors and the endothelium in ADP-mediated relaxation of porcine pancreatic arteries.

Effect of XAC, an adenosine receptor antagonist, and SCH58261, a selective adenosine A_{2A} receptor antagonist

The relaxation to ADP was investigated in the presence of XAC (10 μ M). The relaxation to ADP was largely




Fig. 7 Effect of removal of the endothelium on contraction to UTP in U46619-precontracted porcine pancreatic arteries. Effect of UTP concentration ($F=11.91$, $***P<0.001$) and removal of endothelium reduced the contraction evoked by UTP ($F=43$, $***P<0.001$, two-way ANOVA; $n=10-12$). Data are presented as mean \pm SEM




Fig. 9 Effect of **a** MRS2179 (10 μ M) and **b** the removal of the endothelium on relaxation to ADP in U46619-precontracted porcine pancreatic arteries. With MRS2179 and in endothelium-denuded arteries, effect of ADP concentration ($F=21.42$ and 16.77 , respectively, $***P<0.001$) and MRS2179 and removal of endothelium reduced the contraction evoked by ADP ($F=21.42$ and $F=32.04$, respectively, $***P<0.001$, two-way ANOVA; $n=10-12$). Data are presented as mean \pm SEM




Fig. 10 Effect of XAC (10 μ M), a non-selective adenosine receptor antagonist, and SCH58261 (1 μ M), a selective adenosine A_{2A} receptor antagonist, on relaxation to ADP in U46619-precontracted porcine pancreatic arteries. With XAC and SCH58261, effect of ADP concentrations ($F=7.14$ and $F=6.08$, respectively, $***P<0.001$), XAC and SCH58261 reduced the relaxation evoked by ADP ($F=71.19$ and $F=58.16$, respectively, $***P<0.001$, two-way ANOVA; $n=9-14$). Data are presented as mean \pm SEM

(Fig. 10). This showed that the relaxation to ADP involved A_{2A} adenosine receptors.

Discussion

The current report has provided evidence for the functional expression of contractile P2X₁, P2Y₂ and P2Y₄ receptors and vasorelaxant P2Y₁ and A_{2A} adenosine receptors in porcine pancreatic arteries. These receptors are sensitive to the extracellular nucleotides ATP (P2X₁), UTP (P2Y₂ and P2Y₄) and ADP (P2Y₁ and A_{2A}). The contraction to ATP was endothelium independent, while UTP induced an endothelium-dependent contraction which may involve P2Y₂ and/or P2Y₄ receptors. The relaxation to ADP involved the endothelium and P2Y₁ receptors and A_{2A} adenosine receptors.

A vasoconstrictor response elicited by ATP has been reported in a number of different arteries [21–23]. ATP may also induce vasorelaxation depending on the experimental conditions (level of pre-tone) and relative expression of relevant vasoconstrictile and vasorelaxant receptors [24, 25]. In porcine pancreatic arteries, ATP induced a biphasic response consisting of a contraction followed by a relaxation (Fig. 1b). Since the contraction to ATP was rapidly desensitising, non-cumulative concentration–response curves were investigated. The contractions to ATP and $\alpha\beta$ -meATP were reduced in the presence of suramin, PPADS, $\alpha\beta$ -meATP (a desensitiser of P2X₁ receptors) and NF449 (a P2X₁ selective antagonist) (Figs. 2a, b and 3a, b), which indicates that a large part of the contraction to ATP could be attributed to the activation of P2X₁ receptors. Moreover, the contractile effect of $\alpha\beta$ -meATP is consistent with the expression of P2X₁ receptors in porcine pancreatic arteries. $\alpha\beta$ -meATP was more potent than ATP in eliciting vasoconstriction most likely due

to its greater stability [5]. Since the contraction to ATP was not changed after the endothelium had been removed (Fig. 4a), the expression of P2X₁ receptors was shown to be on the vascular smooth muscle cells (VSMCs). This is consistent with the abundant expression of P2X₁ receptors on VSMCs of most tissues [7].

ATP-induced vasorelaxation was not affected after the endothelium had been removed or in the presence of suramin or PPADS, which suggests that the relaxation to ATP was not due to its action at P2Y receptors. However, the relaxation to ATP was significantly inhibited in the presence of XAC, which suggested an involvement of adenosine receptors expressed on VSMCs of the pancreatic arteries; it is likely that this is due to the activity of adenosine derived from ATP metabolism by ecto-nucleotidase 5'-triphosphate diphosphohydrolase (ENTPDase) enzymes followed by the activity of CD37 and ecto-5'-nucleotidase enzymes [26]. Similarly, in rat coronary arteries, the relaxation to ATP involved P1 receptors, although there was an additional involvement of P2Y receptors [24]. In the current study, further investigation of the adenosine receptor subtypes involved in the relaxation to ATP is required. We and others have shown previously a slow relaxation in response to $\alpha\beta$ -meATP in rat mesenteric arteries, subsequent to contraction [27–29], but we did not observe this in the present study in the porcine pancreatic arteries.

The vasoconstriction to UTP did not desensitise quickly; therefore, cumulative concentration–response curves were used to study the effect of UTP on pancreatic arteries. This contraction was significantly inhibited by suramin and PPADS (Fig. 6), and there was a reduction of the response after the removal of the endothelium (Fig. 7). That would indicate for the first time an endothelium-dependent vasoconstriction evoked by UTP. UTP is known to be active at P2Y₂, P2Y₄ and P2Y₆ receptors [30]. The expression of these receptors in the endothelium and smooth muscle of vessels has been reported [31]. Since MRS2578 was not able to alter the contraction to UTP, this indicates that UTP had no action at P2Y₆ receptors. There are currently no commercially available selective antagonists for either P2Y₂ or P2Y₄ receptors. However, we believe that UTP acted at P2Y₄ receptors since the contraction to UTP was significantly inhibited by both endothelium removal and in the presence of DUP 697, but responses to ATP were unaffected. UTP-induced contraction may also be mediated by P2Y₂ receptors, since MRS2768 which is a selective agonist at P2Y₂ receptors and displays no affinity for P2Y₄ or P2Y₆ receptors was able to evoke a contraction in pancreatic arteries [32] (Fig. 1a).

UTP-induced vasoconstriction has been documented in a number of arteries including rat pulmonary arteries in which the contraction was attributed to P2Y₂ receptors, and in rabbit basilar arteries in which the contraction to UTP was due to action of P2Y₄ receptors [33, 34]. UTP produced an

endothelium-dependent relaxation in rabbit pulmonary arteries and in rat mesenteric arterial bed, but the receptor subtypes were undefined [22, 35]. In bovine middle cerebral arterial strips, UTP had a dual response, and it induced a contraction in endothelium-denuded arteries, but a relaxation in endothelium-intact arteries [36]. The absence of endothelium-dependent or endothelium-independent relaxation to UTP and some other nucleotides in rat renal arteries was reported [37], which is consistent with the current study since there was no evidence of a UTP-mediated relaxation in porcine pancreatic arteries. Hence, porcine pancreatic arteries appear not to express relaxant P2Y₂ and/or P2Y₄ receptors.

To investigate the mechanism underlying the contraction mediated by UTP in pancreatic arteries, the response to UTP was examined in the presence of DUP 697. As seen in Fig. 8, the endothelium-dependent contraction was attenuated in the presence of the selective COX-2 inhibitor. Endothelial cells can release endothelium-derived contractile factors (EDCFs), which may include thromboxane A₂, prostaglandin F_{2α}, leukotrienes and endothelin-1. Thromboxane A₂ and prostaglandin F_{2α} are released from the endothelium due to the activity of COX-2 [38, 39]. The reduction of the contraction to UTP in the presence of DUP 697 indicated the involvement of thromboxane A₂ and prostaglandins in the contraction to UTP. These agents, after being released from the endothelium, may act on their receptors on VSMCs to cause contraction [39]. The different effects of DUP 697 on responses to UTP and ATP further suggest that they are acting on different receptors.

The relaxation to ADP did not desensitise rapidly; therefore, cumulative concentration–response curves were used to study the effect of ADP on pancreatic arteries. The relaxation was significantly attenuated by MRS2179, a selective P2Y₁ receptor antagonist (Fig. 9a). In addition, the relaxation to ADP was reduced after the endothelium had been removed, by a similar extent as observed in the presence of the MRS2179 (Fig. 9b). This may suggest that P2Y₁ receptors are expressed on the endothelium. Indeed, a number of reports show that P2Y₁ receptors are expressed on the endothelium and are responsible for the relaxation of arteries, including rat thoracic aortic and porcine mesenteric arteries [40, 41]. The relaxation to ADP in our study was largely reduced in the presence of XAC and SCH58261 (adenosine receptor antagonists). Adenosine receptors may be expressed on the endothelium or the vascular smooth muscle [42]. Since XAC and SCH58261 produced a greater reduction in the relaxation to ADP than the inhibition induced by removal of the endothelium (Fig. 10), this suggests that relaxation to ADP involves A_{2A} adenosine receptors expressed, at least in part, on VSMCs. The mechanism by which ADP would produce adenosine to act at the adenosine receptors is still to be elucidated. The simplest explanation is that it is broken down by ENTPDases and by CD37 enzymes to adenosine [26].

Alternatively, as suggested in porcine coronary arteries, ADP mediates a relaxation via a mechanism that involves ADP-evoked adenosine release and the subsequent activation of A_{2A} receptors [20]. In contrast to the porcine pancreatic vessels, ADP in rat pancreatic arteries induced a contraction at a high concentration (1 mM); this contraction was similar to that produced by ATP and was much lower than the contraction induced by αβ-meATP [43]. Further investigation is required to determine the involvement of endothelium-derived relaxing factors or endothelium-derived hyperpolarising factors released from the endothelium in the ADP-induced relaxation.

Reduction in pancreatic blood flow has been observed in acute and chronic pancreatitis and some other pancreatic diseases [44, 45], implicating pancreatic tissue perfusion as an important factor in the pathogenesis of pancreatic diseases and symptoms. There is increasing evidence for the role of purinergic signalling in the pathophysiology of the pancreas [2]. Hence, drugs designed to target specific components of purinergic system may be of relevance to the management of pancreatitis, cystic fibrosis, pancreatic cancer and diabetes.

In summary, the functional expression of P2X₁ and A_{2A} adenosine receptors on VSMCs and P2Y₂ and/or P2Y₄ receptors on the endothelium of porcine pancreatic arteries was indicated in the current study. Activation of P2X₁ receptors by ATP or αβ-meATP induced a vasoconstriction, and UTP acts at P2Y₂ and/or P2Y₄ receptors to induce a contraction. ADP and ATP activate A_{2A} adenosine receptors to induce relaxation, together with an action of ADP on P2Y₁ receptors. Pancreatic arteries appear to lack vasorelaxant P2Y₂ and/or P2Y₄ receptors.

Acknowledgments We would like to thank the Damascus University in Syria for funding the project.

References

- Novak I (2008) Purinergic receptors in the endocrine and exocrine pancreas. *Purinergic Signal* 4(3):237–253
- Burnstock G, Novak I (2012) Purinergic signalling in the pancreas in health and disease. *J Endocrinol* 213(2):123–141
- Khakh BS, Burnstock G, Kennedy C, King BF, North RA, Séguéla P et al (2001) International union of pharmacology. XXIV. Current status of the nomenclature and properties of P2X receptors and their subunits. *Pharmacol Rev* 53(1):107–118
- Abbracchio MP, Burnstock G, Boeynaems J-M, Barnard EA, Boyer JL, Kennedy C et al (2006) International union of pharmacology LVIII: update on the P2Y G protein-coupled nucleotide receptors: from molecular mechanisms and pathophysiology to therapy. *Pharmacol Rev* 58(3):281–341
- Kasakov L, Burnstock G (1982) The use of the slowly degradable analog, α,β-methylene ATP, to produce desensitisation of the P2-purinoceptor: effect on non-adrenergic, non-cholinergic responses of the guinea-pig urinary bladder. *Eur J Pharmacol* 86(2):291–294

- 538 6. Burnstock G (2006) Pathophysiology and therapeutic potential of
539 purinergic signaling. *Pharmacol Rev* 58(1):58–86
- 540 7. Kügelgen I (2008) Pharmacology of mammalian P2X and P2Y
541 receptors. *Biotrend Reviews*, no. 3
- 542 8. Communi D, Govaerts C, Parmentier M, Boeynaems J-M (1997)
543 Cloning of a human purinergic P2Y receptor coupled to phospholi-
544 pase C and adenylyl cyclase. *J Biol Chem* 272(51):31969–31973
- 545 9. Qi A-D, Zamboni AC, Insel PA, Nicholas RA (2001) An arginine/
546 glutamine difference at the juxtaposition of transmembrane domain 6
547 and the third extracellular loop contributes to the markedly different
548 nucleotide selectivities of human and canine P2Y₁₁ receptors. *Mol*
549 *Pharmacol* 60(6):1375–1382
- 550 10. Léon C, Hechler B, Vial C, Leray C, Cazenave J-P, Gachet C (1997) The
551 P2Y₁ receptor is an ADP receptor antagonized by ATP and expressed in
552 platelets and megakaryoblastic cells. *FEBS Lett* 403(1):26–30
- 553 11. Bodor ET, Waldo GL, Hooks SB, Corbitt J, Boyer JL, Harden TK
554 (2003) Purification and functional reconstitution of the human P2Y₁₂
555 receptor. *Mol Pharmacol* 64(5):1210–1216
- 556 12. Marteau F, Le Poul E, Communi D, Communi D, Labouret C, Savi P
557 et al (2003) Pharmacological characterization of the human P2Y₁₃
558 receptor. *Mol Pharmacol* 64(1):104–112
- 559 13. Waldo GL, Harden TK (2004) Agonist binding and Gq-stimulating
560 activities of the purified human P2Y₁ receptor. *Mol Pharmacol* 65(2):
561 426–436
- 562 14. Nicholas RA, Lazarowski ER, Watt WC, Li Q, Boyer J, Harden TK
563 (1996) Pharmacological and second messenger signalling selectiv-
564 ities of cloned P2Y receptors. *J Auton Pharmacol* 16(6):319–323
- 565 15. Carter RL, Fricks IP, Barrett MO, Burianek LE, Zhou Y, Ko H et al
566 (2009) Quantification of G_i-mediated inhibition of adenylyl cyclase
567 activity reveals that UDP is a potent agonist of the human P2Y₁₄
568 receptor. *Mol Pharmacol* 76(6):1341–1348
- 569 16. Coutinho-Silva R, Parsons M, Robson T, Lincoln J, Burnstock G
570 (2003) P2X and P2Y purinoceptor expression in pancreas from
571 streptozotocin-diabetic rats. *Mol Cell Endocrinol* 204(1–2):141–154
- 572 17. Hillaire-Buys D, Chapal J, Petit P, Loubatières-Mariani M-M (1991)
573 Dual regulation of pancreatic vascular tone by P2X and P2Y
574 purinoceptor subtypes. *Eur J Pharmacol* 199(3):309–314
- 575 18. Hillaire-Buys D, Dietz S, Chapal J, Petit P, Loubatières-Mariani MM
576 (1999) Involvement of P2X and P2U receptors in the constrictor
577 effect of ATP on the pancreatic vascular bed. *Journal de la Societe*
578 *de Biologie* 193(1):57–61
- 579 19. Alsaqati M, Chan SLF, Ralevic R (2011) Characterisation of the
580 response to ADP in porcine isolated pancreatic arteries. [http://www.](http://www.pa2online.org/abstracts/vol8issue1abst028p.pdf)
581 [pa2online.org/abstracts/vol8issue1abst028p.pdf](http://www.pa2online.org/abstracts/vol8issue1abst028p.pdf)
- 582 20. Rayment SJ, Ralevic V, Barrett DA, Cordell R, Alexander SPH
583 (2007) A novel mechanism of vasoregulation: ADP-induced relaxa-
584 tion of the porcine isolated coronary artery is mediated via adenosine
585 release. *FASEB J* 21(2):577–585
- 586 21. Kügelgen I, Häussinger D, Starker K (1987) Evidence for a
587 vasoconstriction-mediating receptor for UTP, distinct from the P2
588 purinoceptor, in rabbit ear artery. *Naunyn Schmiedebergs Arch*
589 *Pharmacol* 336(5):556–560
- 590 22. Ralevic V, Burnstock G (1991) Effects of purines and pyrimidines on
591 the rat mesenteric arterial bed. *Circ Res* 69(6):1583–1590
- 592 23. Rubino A, Burnstock G (1996) Evidence for a P2-purinoceptor
593 mediating vasoconstriction by UTP, ATP and related nucleotides in
594 the isolated pulmonary vascular bed of the rat. *Br J Pharmacol*
595 118(6):1415–1420
- 596 24. Korchazhkina O, Wright G, Exley C (1999) Intravascular ATP and
597 coronary vasodilation in the isolated working rat heart. *Br J*
598 *Pharmacol* 127(3):701–708
- 599 25. Ralevic V, Burnstock G (1996) Relative contribution of P2U- and
600 P2Y-purinoceptors to endothelium-dependent vasodilatation in the
601 golden hamster isolated mesenteric arterial bed. *Br J Pharmacol*
602 117(8):1797–1802
- 603 26. Zimmermann H (2000) Extracellular metabolism of ATP and other
604 nucleotides. *Naunyn Schmiedebergs Arch Pharmacol* 362(4–5):299–
605 309
- 606 27. Stanford SJ, Mitchell JA (1998) ATP-induced vasodilatation in the
607 rat isolated mesenteric bed exhibits two apparent phases. *Br J*
608 *Pharmacol* 125:94P
- 609 28. Ralevic V (2001) Mechanism of prolonged vasorelaxation to ATP in
610 the rat isolated mesenteric arterial bed. *Br J Pharmacol* 132(3):685–
611 692
- 612 29. Ralevic V (2002) The involvement of smooth muscle P2X
613 receptors in the prolonged vasorelaxation response to purine
614 nucleotides in the rat mesenteric arterial bed. *Br J Pharmacol*
615 135(8):1988–1994
- 616 30. Burnstock G, Williams M (2000) P2 purinergic receptors: modula-
617 tion of cell function and therapeutic potential. *J Pharmacol Exp Ther*
618 295(3):862–869
- 619 31. Burnstock G (2007) Purine and pyrimidine receptors. *CMLS* 64(12):
620 1471–1483
- 621 32. Ko H, Carter RL, Cosyn L, Petrelli R, de Castro S, Besada P et al
622 (2008) Synthesis and potency of novel uracil nucleotides and deriv-
623 atives as P2Y₂ and P2Y₆ receptor agonists. *Bioorg Med Chem*
624 16(12):6319–6332
- 625 33. Hartley SA, Kato K, Salter KJ, Kozlowski RZ (1998) Functional
626 evidence for a novel suramin-insensitive pyrimidine receptor in rat
627 small pulmonary arteries. *Circ Res* 83(9):940–946
- 628 34. Miyagi Y, Zhang JH (2004) α , β -Methylene ATP enhances P2Y₄
629 contraction of rabbit basilar artery. *Am J Physiol Heart Circ Physiol*
630 286(4):H1546–H1551
- 631 35. Qasabian RA, Schyvens C, Owe-Young R, Killen JP, Macdonald PS,
632 Conigrave AD et al (1997) Characterization of the P2 receptors in
633 rabbit pulmonary artery. *Br J Pharmacol* 120(4):553–558
- 634 36. Miyagi Y, Kobayashi S, Nishimura J, Fukui M, Kanaide H (1996)
635 Dual regulation of cerebrovascular tone by UTP: P2U receptor-
636 mediated contraction and endothelium-dependent relaxation. *Br J*
637 *Pharmacol* 118(4):847–856
- 638 37. Knight GE, Oliver-Redgate R, Burnstock G (2003) Unusual absence
639 of endothelium-dependent or -independent vasodilatation to purines
640 or pyrimidines in the rat renal artery. *Kidney Int* 64(4):1389–1397
- 641 38. Mombouli JV, Vanhoutte PM (1993) Purinergic endothelium-
642 dependent and -independent contractions in rat aorta. *Hypertension*
643 22(4):577–583
- 644 39. Wong SL, Leung FP, Lau CW, Au CL, Yung LM, Yao X et al (2009)
645 Cyclooxygenase-2-derived prostaglandin F₂ α mediates
646 endothelium-dependent contractions in the aortae of hamsters with
647 increased impact during aging. *Circ Res* 104(2):228–235
- 648 40. Dol-Gleizes F, Mares AM, Savi P, Herbert JM (1999) Relaxant effect
649 of 2-methyl-thio-adenosine diphosphate on rat thoracic aorta: effect
650 of clopidogrel. *Eur J Pharmacol* 367(2–3):247–253
- 651 41. Alefishat E, Alexander S, Ralevic V (2010) Effect of palmitoyl CoA
652 on ADP-evoked vasorelaxations in porcine isolated coronary and
653 mesenteric arteries. [http://www.fasebj.org/cgi/content/meeting_](http://www.fasebj.org/cgi/content/meeting_abstract/24/1_MeetingAbstracts/lb426)
654 [abstract/24/1_MeetingAbstracts/lb426](http://www.fasebj.org/cgi/content/meeting_abstract/24/1_MeetingAbstracts/lb426)
- 655 42. Schulte G, Fredholm BB (2003) Signalling from adenosine receptors
656 to mitogen-activated protein kinases. *Cell Signal* 15(9):813–827
- 657 43. Chapal J, Loubatières-Mariani MM (1983) Evidence for purinergic
658 receptors on vascular smooth muscle in rat pancreas. *Eur J Pharmacol*
659 87(4):423–430
- 660 44. Satoh A, Shimosegawa T, Satoh K, Ito H, Kohno Y, Masamune A
661 et al (2000) Activation of adenosine A₁-receptor pathway induces
662 edema formation in the pancreas of rats. *Gastroenterology* 119(3):
663 829–836
- 664 45. Nguyen NC, Taalab K, Osman MM (2010) Decreased blood flow
665 with increased metabolic activity: a novel sign of pancreatic tumor
666 aggressiveness. *Clin Cancer Res Off J Am Assoc Cancer Res* 16(1):
667 367, author reply 567

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES.

- Q1. Kindly check the inserted spelled out form 'standard error of the mean' of the abbreviation 'SEM' if correct.
- Q2. Kindly check the inserted spelled out form 'half maximal effective concentration' of the abbreviation 'EC₅₀' if correct.
- Q3. Kindly provide access date for references 'Alsaqati et al. [19]' and 'Alefishat et al. [41]' in the list.

UNCORRECTED PROOF